

Anti-Cldn1 antibody



Catalog Number: 175334

Product name

Anti-Cldn1 antibody

Specificity

Human, Mouse, Rat

Antibody description

Rabbit polyclonal antibody to Cldn1

Preparation

This antigen of this antibody was klh conjugated synthetic peptide derived from mouse claudin 1 121-211/211

Formulation

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Storage

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

Clonality

Polyclonal

Ig Type

Rabbit IgG

Applications

WB, IHC-P, FC

Dilutions

WB:1:500-2000

IHC-P:1:400-800

FC:1µg/Test

Validations




Fig1: Blank control: Raji(blue).; Primary Antibody:Rabbit Anti-Claudin 1 antibody , Dilution: 1µg in 100 µL 1X PBS containing 0.5% BSA;; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions);; Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.; Protocol; The cells were fixed with 2% paraformaldehyde (10 min). Antibody (1µg /1x10⁶ cells) were incubated for 30 min on the ice, followed by 1 X PBS containing 0.5% BSA + 1 0% goat serum (15 min) to block non-specific protein-protein interactions. Then the Goat Anti-rabbit IgG/PE antibody was added into the blocking buffer mentioned above to react with the primary antibody of 175334# at 1/200 dilution for 30 min on ice. Acquisition of 20,000 events was performed.




Fig2: Tissue/cell: rat intestine tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;; Antigen retrieval: citrate buffer (0.01M, pH 6.0),

Anti-Cldn1 antibody



Catalog Number: 175334

Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;; Incubation: Anti-Claudin-1 Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010)




Fig3: Sample:; HepG2(Human) Cell Lysate at 30 ug; Primary: Anti-Claudin 1 at 1/1000 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 23 kD;




Fig4: Sample:; Kidney (Mouse) Lysate at 40 ug; Primary: Anti-Claudin 1 at 1/1000 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 23 kD;




Fig5: Sample:; A431(Human) Cell Lysate at 30 ug; Primary: Anti-Claudin 1 at 1/1000 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 23 kD; Observed band size: 23 kD