

Human CD16a/FCGR3A (His & AVI Tag) recombinant protein



Catalog Number: 503735

General Information

Gene Name Synonym

CD16a antigen; Fc-gamma RIII-alpha; FcR-10; IgG Fc receptor III-2

Protein Construction

A DNA sequence encoding the extracellular domain (Met 1-Gln 208) of human CD16a (P08637-1) was fused with a c-terminal polyhistidine tagged AVI tag at the C-terminus. It is identical to FCGR3A158F/V in the reference.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Using the Octet RED System, the affinity constant (Kd) of CD16a bound to Human IgG1 was 30nM.

Purity

> 95 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μ g of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Gly 17

Molecular Mass

The secreted recombinant human CD16a consists of 223 amino acids and predicts a molecular mass of 25.1 kDa. By SDS-PAGE under reducing conditions, the apparent molecular mass of the protein is approximately 46 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.2

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human CD16a / FCGR3A Protein (176 Phe) (His & AVI Tag) SDS-PAGE