Human B7-DC/PD-L2/CD273 (Fc Tag) recombinant protein

Catalog Number: 500765



General Information

Gene Name Synonym

Butyrophilin B7-DC

Protein Construction

A DNA sequence encoding the human PDCD1LG2 (NP_079515.2) (Met1-Pro219) was expressed with the Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

- 1. Measured by its binding ability in a functional ELISA.
- 2. Immobilized human PD1-His (Cat:503606) at 10 μ g/mL (100 μ L/well) can bind human PDCD1LG2-Fc. The EC₅₀ of PDCD1LG2-Fc is 0.05-0.12 μ g/mL.

Purity

> 95 % as determined by SDS-PAGE.

Endotoxin

<1.0 EU per μg protein as determined by the LAL method.

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Leu 20

Molecular Mass

The recombinant human PDCD1LG2 consists 438 amino acids and predicts a molecular mass of 49.3 kDa.

Formulation

Lyophilized from sterile PBS, pH 7.4.

- $1.\,5$ % trehalose and mannitol are added as protectants before lyophilization.
- 2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human B7-DC / PD-L2 / CD273 / PDCD1LG2 Protein (Fc Tag) SDS-PAGE