

General Information

Protein Construction

A DNA sequence encoding the mouse IFNG (P01580) (Met 1-Cys 155) was fused with the Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Activity

1. Measured in antiviral assays using L929 cells infected with vesicular stomatitisvirus (VSV). The ED₅₀ for this effect is typically 20-80 ng/mL.
2. Measured by its ability to bind with recombinant mouse IFNGR1-His (Cat:502661) in a functional ELISA.

Purity

> 92 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

His 23

Molecular Mass

The secreted recombinant mouse IFNG/Fc is a disulfide-linked homodimer. The reduced monomer comprises 374 amino acids and has a calculated molecular mass of 42.5 kDa. As a result of glycosylation, the apparent molecular mass of the monomer is approximately 55 kDa in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse IFNG / Interferon Gamma Protein (Fc Tag)
SDS-PAGE