# **Anti-DSC2 antibody**

Catalog Number: 103536



#### **Product name**

Anti-DSC2 antibody

#### **Immunogen**

Human DSC2 (His Tag) recombinant protein

# **Specificity**

Human DSC2

# **Antibody description**

Rabbit polyclonal to DSC2

# **Preparation**

Produced in rabbits immunized with purified, recombinant Human DSC2 (rh DSC2; Q02487-1; Met 1-Arg 684). DSC2 specific IgG was purified by Human DSC2 affinity chromatography.

#### **Formulation**

0.2 µm filtered solution in PBS with 5% trehalose

#### **Storage**

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

### **Clonality**

Polyclonal

#### Ig Type

Rabbit IgG

## **Applications**

ELISA, IHC-P


#### **Dilutions**

ELISA: 0.1-0.2 μg/mL

This antibody can be used at 0.1-0.2  $\mu$ g/mL with the appropriate secondary reagents to detect Human DSC2. The detection limit for Human DSC2 is approximately 0.00245 ng/well.


IHC-P:  $0.1-2 \mu g/mL$ 

#### **Validations**



DSC2 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human DSC2 in human esophagus with rabbit polyclonal antibody (1  $\mu$ g/mL, formalin-fixed paraffin embedded sections). Positive staining was localized to membrane of the squamous epithelium.



DSC2 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human DSC2 in human heart with rabbit polyclonal antibody (1

# **Anti-DSC2 antibody**

Catalog Number: 103536



 $\mu g/mL$ , formalin-fixed paraffin embedded sections). Positive staining was localized to intercalated disk. The left panel: tissue incubated

with primary antibody; The right panel: tissue incubated with the mixture of primary antibody and antigen (recombinant protein).